

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(3-Methoxybenzoyl)-2-methylbenzenesulfonamide

S. Sreenivasa,^a D. Darshan,^b M. Prakash Shet,^c
N. R. Mohan,^a Vijith Kumar^d and P. A. Suchetan^{c*}^aDepartment of Studies and Research in Chemistry, Tumkur University, Tumkur, Karnataka 572 103, India, ^bUniversity College of Science, Tumkur University, Tumkur, India, ^cDepartment of Studies and Research in Chemistry, U.C.S., Tumkur University, Tumkur, Karnataka 572 103, India, and ^dSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, India
Correspondence e-mail: pasuchetan@yahoo.co.in


Received 29 June 2013; accepted 4 July 2013

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}—\text{C}) = 0.004$ Å; R factor = 0.045; wR factor = 0.118; data-to-parameter ratio = 13.1.

In the title compound, $\text{C}_{15}\text{H}_{15}\text{NO}_4\text{S}$, the dihedral angle between the methyl- and methoxy-substituted benzene rings is 88.99 (12)°. An intramolecular $\text{C}—\text{H} \cdots \text{O}$ hydrogen bond occurs. In the crystal, adjacent molecules form inversion-related dimers through strong $\text{N}—\text{H} \cdots \text{O}$ hydrogen bonds, generating $R_2^2(8)$ loops. The dimers are further connected through $\text{C}—\text{H} \cdots \text{O}$ interactions that form $C(8)$ chains parallel to (001). Molecules are also connected through other $\text{C}—\text{H} \cdots \text{O}$ hydrogen bonds along the b axis, forming additional $C(8)$ chains. Two aromatic $\pi—\pi$ stacking interactions [centroid-centroid separations = 3.6150 (1) and 3.6837 (1) Å] generate a three-dimensional architecture.

Related literature

For similar structures, see: Gowda *et al.* (2010); Suchetan *et al.* (2010, 2011).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{15}\text{NO}_4\text{S}$
 $M_r = 305.34$
Monoclinic, $C2/c$ $a = 26.713$ (5) Å
 $b = 7.3717$ (4) Å
 $c = 19.636$ (3) Å $\beta = 131.21$ (3)°
 $V = 2908.7$ (7) Å³
 $Z = 8$
Mo $K\alpha$ radiation $\mu = 0.24$ mm⁻¹
 $T = 293$ K
 $0.33 \times 0.27 \times 0.22$ mm

Data collection

Bruker APEXII CCD area-detector
diffractometer
5294 measured reflections2558 independent reflections
1970 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.118$
 $S = 1.04$
2558 reflections
196 parametersH atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³**Table 1**
Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
$\text{N1}—\text{H}\text{N1} \cdots \text{O2}^{\text{i}}$	0.80 (3)	2.15 (4)	2.929 (4)	166
$\text{C3}—\text{H3} \cdots \text{O3}^{\text{ii}}$	0.93	2.60	3.463 (3)	155
$\text{C10}—\text{H10} \cdots \text{O2}^{\text{i}}$	0.93	2.60	3.404 (3)	145
$\text{C15}—\text{H15B} \cdots \text{O3}^{\text{iii}}$	0.96	2.36	3.323 (3)	178
$\text{C6}—\text{H6} \cdots \text{O1}$	0.93	2.46	2.861 (4)	106

Symmetry codes: (i) $-x, y, -z + \frac{1}{2}$; (ii) $x, y - 1, z$; (iii) $x, -y + 2, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: APEX2 and SAINT-Plus (Bruker, 2009); data reduction: SAINT-Plus and XPREP (Bruker, 2009); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXL97.

The authors thank Professor T. N. Guru Row, Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, for his help and valuable suggestions. PAS and MPS thank the University Grants Commission (UGC), India, for financial support under its Minor Research Project scheme.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5340).

References

- Bruker (2009). APEX2, SADABS, SAINT-Plus and XPREP. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2010). *Acta Cryst.* **E66**, o433.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Suchetan, P. A., Foro, S. & Gowda, B. T. (2011). *Acta Cryst.* **E67**, o3489.
- Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2010). *Acta Cryst.* **E66**, o1024.

supplementary materials

Acta Cryst. (2013). E69, o1232 [doi:10.1107/S1600536813018539]

***N*-(3-Methoxybenzoyl)-2-methylbenzenesulfonamide**

S. Sreenivasa, D Darshan, M. Prakash Shet, N. R. Mohan, Vijith Kumar and P. A. Suchetan

Comment

As a part of our continued efforts to study the crystal structures of *N*-(aroyl)-arylsulfonamides (Suchetan *et al.*, 2010, 2011), we report herein the crystal structure of the title compound (I).

In the title compound, C₁₅H₁₅NO₄S, the dihedral angle between the benzene rings is 88.99°. In the molecule, the conformation between the N-H bond and the ortho-methyl group in the sulfonyl benzene ring is *syn*. This is similar to what is observed in *N*-(benzoyl)-2-methylbenzenesulfonamide (II, Suchetan *et al.*, 2010), *N*-(3-chlorobenzoyl)-2-methylbenzenesulfonamide (III, Suchetan *et al.*, 2011) and *N*-(3-methylbenzoyl)-2-methylbenzenesulfonamide (IV, Gowda *et al.*, 2010). Similarly, the conformation between the N-H bond and the meta-methoxy group in the benzoyl ring is *syn*, also similar to the conformation in III (Suchetan *et al.*, 2011) and IV (Gowda *et al.*, 2010).

Adjacent molecules form inversion related dimers through strong N1—HN1···O2 hydrogen bonds, Table 1, generating R₂²(8) loops. The dimers are further connected through intermolecular C10—H10···O2 interactions that form C(8) chains parallel to (001). Molecules are also connected through other intermolecular C15—H15B···O3 hydrogen bonds along the *b* axis forming additional C(8) chains. Two aromatic π – π stacking interactions (centroid-centroid separations 3.6150 (1) Å and 3.6837 (1) Å) are also observed.

Experimental

The title compound was prepared by refluxing a mixture of 3-methoxybenzoic acid, 2-methylbenzenesulfonamide and phosphorus oxychloride, POCl₃, for 2 h on a water bath. The resultant mixture was cooled and poured into ice cold water. The solid obtained was filtered and washed thoroughly with water and then dissolved in sodium bicarbonate solution. The compound was later reprecipitated by acidifying the filtered solution with dilute HCl. The filtered and dried solid was recrystallized to the constant melting point (423 K).

Colorless prisms of (I) were obtained from a slow evaporation of its ethanolic solution at room temperature.

Refinement

The H atom of the NH group was located in a difference map and later refined freely. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the *U*_{eq} of the parent atom).

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *APEX2* and *SAINT-Plus* (Bruker, 2009); data reduction: *SAINT-Plus* and *XPREF* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).




Figure 1

Molecular structure of the title compound, showing the atom-labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Figure 2

Molecular packing of (I) with hydrogen bonding shown as dashed lines. Hydrogen atoms bound to carbon are omitted for clarity.




Figure 3

Display of C—H...O interactions among molecules along *b* axis forming C(8) chains.




Figure 4

Stacking of molecules through *Cg*...*Cg* interactions. *Cg*1 and *Cg*2 are the centroids of the carbonyl and sulfonyl bounded benzene rings respectively.

***N*-(3-Methoxybenzoyl)-2-methylbenzenesulfonamide**

Crystal data

$C_{15}H_{15}NO_4S$

$M_r = 305.34$

Monoclinic, *C2/c*

Hall symbol: $-C\ 2yc$

$a = 26.713\ (5)\ \text{\AA}$

$b = 7.3717\ (4)\ \text{\AA}$

$c = 19.636\ (3)\ \text{\AA}$

$\beta = 131.21\ (3)^\circ$

$V = 2908.7\ (7)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1280$

Prism

$D_x = 1.394\ \text{Mg m}^{-3}$

Melting point: 423 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1232 reflections

$\theta = 2.8\text{--}25.0^\circ$

$\mu = 0.24\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colourless

$0.33 \times 0.27 \times 0.22\ \text{mm}$

Data collection

Bruker APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 5294 measured reflections
 2558 independent reflections

1970 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$
 $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.8^\circ$
 $h = -31 \rightarrow 31$
 $k = -8 \rightarrow 8$
 $l = -14 \rightarrow 23$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.118$
 $S = 1.04$
 2558 reflections
 196 parameters
 0 restraints
 0 constraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0544P)^2 + 0.0171P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
HN1	0.0737 (12)	0.796 (3)	0.2976 (17)	0.055 (8)*
C1	0.05769 (11)	0.6312 (3)	0.13024 (14)	0.0444 (5)
C9	0.18751 (12)	0.9525 (3)	0.40254 (15)	0.0425 (5)
C10	0.16739 (12)	0.9650 (3)	0.45194 (16)	0.0479 (6)
H10	0.1231	0.9451	0.4237	0.057*
C8	0.14165 (12)	0.9088 (3)	0.30428 (16)	0.0467 (6)
C2	0.06638 (11)	0.4482 (3)	0.15570 (16)	0.0488 (6)
C6	0.07526 (12)	0.6977 (3)	0.08233 (16)	0.0546 (6)
H6	0.0689	0.8195	0.0662	0.066*
C14	0.25399 (13)	0.9826 (3)	0.44521 (17)	0.0529 (6)
H14	0.2678	0.9748	0.4125	0.063*
C11	0.21314 (12)	1.0071 (3)	0.54308 (17)	0.0496 (6)
C13	0.29856 (13)	1.0235 (3)	0.53520 (18)	0.0602 (7)
H13	0.3429	1.0430	0.5635	0.072*
C12	0.27921 (12)	1.0365 (3)	0.58542 (17)	0.0529 (6)
H12	0.3102	1.0646	0.6468	0.063*
C5	0.10253 (13)	0.5803 (4)	0.05870 (18)	0.0653 (7)

H5	0.1146	0.6233	0.0267	0.078*
C4	0.11146 (14)	0.4011 (4)	0.08281 (18)	0.0666 (8)
H4	0.1296	0.3227	0.0670	0.080*
C3	0.09384 (13)	0.3358 (3)	0.13017 (18)	0.0601 (7)
H3	0.1004	0.2136	0.1457	0.072*
C7	0.04728 (14)	0.3666 (3)	0.20628 (19)	0.0651 (7)
H7A	0.0607	0.4466	0.2543	0.098*
H7B	0.0691	0.2516	0.2312	0.098*
H7C	−0.0001	0.3496	0.1655	0.098*
C15	0.23539 (16)	1.0486 (5)	0.68140 (19)	0.0927 (11)
H15A	0.2674	0.9521	0.7117	0.139*
H15B	0.2121	1.0546	0.7032	0.139*
H15C	0.2578	1.1616	0.6934	0.139*
O1	0.00755 (9)	0.9515 (2)	0.10733 (11)	0.0614 (5)
O2	−0.02956 (7)	0.7037 (2)	0.14851 (11)	0.0527 (4)
O3	0.15524 (9)	0.9414 (2)	0.25752 (12)	0.0646 (5)
O4	0.18890 (10)	1.0151 (3)	0.58597 (13)	0.0771 (6)
S1	0.02251 (3)	0.78797 (8)	0.15606 (4)	0.0461 (2)
N1	0.08109 (10)	0.8300 (3)	0.26647 (13)	0.0456 (5)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0325 (12)	0.0577 (14)	0.0369 (12)	−0.0007 (11)	0.0202 (11)	−0.0079 (10)
C9	0.0428 (13)	0.0396 (11)	0.0460 (14)	−0.0036 (10)	0.0296 (12)	−0.0046 (10)
C10	0.0408 (14)	0.0560 (13)	0.0514 (15)	−0.0117 (11)	0.0323 (13)	−0.0128 (11)
C8	0.0477 (15)	0.0498 (13)	0.0500 (15)	−0.0079 (11)	0.0354 (14)	−0.0059 (11)
C2	0.0370 (13)	0.0540 (13)	0.0470 (14)	−0.0035 (11)	0.0240 (13)	−0.0114 (11)
C6	0.0477 (15)	0.0703 (16)	0.0468 (15)	0.0002 (12)	0.0315 (14)	−0.0041 (12)
C14	0.0455 (15)	0.0674 (15)	0.0538 (16)	−0.0046 (12)	0.0361 (14)	−0.0031 (12)
C11	0.0479 (15)	0.0570 (13)	0.0509 (15)	−0.0084 (12)	0.0355 (14)	−0.0110 (11)
C13	0.0386 (14)	0.0809 (17)	0.0573 (17)	−0.0073 (13)	0.0299 (14)	−0.0053 (13)
C12	0.0424 (15)	0.0643 (15)	0.0435 (15)	−0.0053 (12)	0.0247 (13)	−0.0060 (11)
C5	0.0528 (17)	0.100 (2)	0.0516 (16)	−0.0027 (16)	0.0381 (15)	−0.0121 (15)
C4	0.0513 (17)	0.087 (2)	0.0610 (18)	0.0025 (15)	0.0369 (16)	−0.0217 (15)
C3	0.0491 (16)	0.0606 (15)	0.0614 (17)	0.0022 (13)	0.0324 (15)	−0.0135 (13)
C7	0.0658 (19)	0.0565 (15)	0.075 (2)	−0.0007 (13)	0.0473 (18)	0.0007 (13)
C15	0.071 (2)	0.164 (3)	0.0525 (19)	−0.016 (2)	0.0442 (19)	−0.0258 (19)
O1	0.0668 (12)	0.0571 (9)	0.0569 (12)	0.0121 (9)	0.0393 (11)	0.0084 (8)
O2	0.0360 (9)	0.0696 (10)	0.0542 (10)	−0.0035 (8)	0.0305 (9)	−0.0118 (8)
O3	0.0606 (12)	0.0922 (13)	0.0561 (12)	−0.0179 (10)	0.0449 (11)	−0.0102 (9)
O4	0.0531 (12)	0.1345 (16)	0.0517 (12)	−0.0218 (11)	0.0380 (11)	−0.0281 (11)
S1	0.0400 (4)	0.0541 (4)	0.0427 (4)	0.0021 (3)	0.0267 (3)	−0.0040 (3)
N1	0.0450 (12)	0.0577 (12)	0.0394 (12)	−0.0091 (9)	0.0301 (11)	−0.0074 (9)

Geometric parameters (Å, °)

C1—C6	1.388 (3)	C13—H13	0.9300
C1—C2	1.404 (3)	C12—H12	0.9300
C1—S1	1.762 (2)	C5—C4	1.370 (4)

C9—C10	1.388 (3)	C5—H5	0.9300
C9—C14	1.394 (3)	C4—C3	1.376 (4)
C9—C8	1.487 (3)	C4—H4	0.9300
C10—C11	1.382 (3)	C3—H3	0.9300
C10—H10	0.9300	C7—H7A	0.9600
C8—O3	1.213 (3)	C7—H7B	0.9600
C8—N1	1.387 (3)	C7—H7C	0.9600
C2—C3	1.400 (3)	C15—O4	1.431 (3)
C2—C7	1.509 (3)	C15—H15A	0.9600
C6—C5	1.393 (3)	C15—H15B	0.9600
C6—H6	0.9300	C15—H15C	0.9600
C14—C13	1.363 (3)	O1—S1	1.4232 (16)
C14—H14	0.9300	O2—S1	1.4391 (16)
C11—O4	1.360 (3)	S1—N1	1.666 (2)
C11—C12	1.385 (3)	N1—HN1	0.79 (2)
C13—C12	1.386 (3)		
C6—C1—C2	122.0 (2)	C4—C5—H5	120.2
C6—C1—S1	116.49 (18)	C6—C5—H5	120.2
C2—C1—S1	121.48 (17)	C5—C4—C3	120.7 (2)
C10—C9—C14	119.7 (2)	C5—C4—H4	119.6
C10—C9—C8	123.6 (2)	C3—C4—H4	119.6
C14—C9—C8	116.6 (2)	C4—C3—C2	121.9 (2)
C11—C10—C9	120.0 (2)	C4—C3—H3	119.1
C11—C10—H10	120.0	C2—C3—H3	119.1
C9—C10—H10	120.0	C2—C7—H7A	109.5
O3—C8—N1	120.2 (2)	C2—C7—H7B	109.5
O3—C8—C9	122.5 (2)	H7A—C7—H7B	109.5
N1—C8—C9	117.3 (2)	C2—C7—H7C	109.5
C3—C2—C1	116.3 (2)	H7A—C7—H7C	109.5
C3—C2—C7	119.0 (2)	H7B—C7—H7C	109.5
C1—C2—C7	124.7 (2)	O4—C15—H15A	109.5
C1—C6—C5	119.4 (2)	O4—C15—H15B	109.5
C1—C6—H6	120.3	H15A—C15—H15B	109.5
C5—C6—H6	120.3	O4—C15—H15C	109.5
C13—C14—C9	119.5 (2)	H15A—C15—H15C	109.5
C13—C14—H14	120.2	H15B—C15—H15C	109.5
C9—C14—H14	120.2	C11—O4—C15	117.5 (2)
O4—C11—C10	115.8 (2)	O1—S1—O2	118.18 (11)
O4—C11—C12	124.0 (2)	O1—S1—N1	108.93 (10)
C10—C11—C12	120.2 (2)	O2—S1—N1	103.29 (10)
C14—C13—C12	121.4 (2)	O1—S1—C1	109.21 (11)
C14—C13—H13	119.3	O2—S1—C1	110.47 (10)
C12—C13—H13	119.3	N1—S1—C1	105.95 (11)
C11—C12—C13	119.1 (2)	C8—N1—S1	122.61 (17)
C11—C12—H12	120.4	C8—N1—HN1	120.7 (19)
C13—C12—H12	120.4	S1—N1—HN1	116.5 (19)
C4—C5—C6	119.6 (2)		

C14—C9—C10—C11	0.0 (3)	C5—C4—C3—C2	−0.1 (4)
C8—C9—C10—C11	−179.7 (2)	C1—C2—C3—C4	0.1 (4)
C10—C9—C8—O3	161.3 (2)	C7—C2—C3—C4	179.1 (2)
C14—C9—C8—O3	−18.4 (3)	C10—C11—O4—C15	176.7 (2)
C10—C9—C8—N1	−17.9 (3)	C12—C11—O4—C15	−2.7 (4)
C14—C9—C8—N1	162.4 (2)	O2—O2—S1—O1	0.00 (5)
C6—C1—C2—C3	0.0 (3)	O2—O2—S1—N1	0.00 (6)
S1—C1—C2—C3	178.99 (18)	O2—O2—S1—C1	0.00 (4)
C6—C1—C2—C7	−178.9 (2)	C6—C1—S1—O1	10.5 (2)
S1—C1—C2—C7	0.1 (3)	C2—C1—S1—O1	−168.48 (18)
C2—C1—C6—C5	−0.2 (4)	C6—C1—S1—O2	142.13 (18)
S1—C1—C6—C5	−179.17 (18)	C2—C1—S1—O2	−36.9 (2)
C10—C9—C14—C13	0.2 (3)	C6—C1—S1—O2	142.13 (18)
C8—C9—C14—C13	179.9 (2)	C2—C1—S1—O2	−36.9 (2)
C9—C10—C11—O4	−179.6 (2)	C6—C1—S1—N1	−106.65 (19)
C9—C10—C11—C12	−0.1 (3)	C2—C1—S1—N1	74.3 (2)
C9—C14—C13—C12	−0.2 (4)	O3—C8—N1—S1	−3.9 (3)
O4—C11—C12—C13	179.5 (2)	C9—C8—N1—S1	175.28 (15)
C10—C11—C12—C13	0.1 (4)	O1—S1—N1—C8	−54.9 (2)
C14—C13—C12—C11	0.1 (4)	O2—S1—N1—C8	178.66 (18)
C1—C6—C5—C4	0.2 (4)	O2—S1—N1—C8	178.66 (18)
C6—C5—C4—C3	0.0 (4)	C1—S1—N1—C8	62.5 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N1—Hm1 \cdots O2 ⁱ	0.80 (3)	2.15 (4)	2.929 (4)	166
C3—H3 \cdots O3 ⁱⁱ	0.93	2.60	3.463 (3)	155
C10—H10 \cdots O2 ⁱ	0.93	2.60	3.404 (3)	145
C15—H15B \cdots O3 ⁱⁱⁱ	0.96	2.36	3.323 (3)	178
C6—H6 \cdots O1	0.93	2.46	2.861 (4)	106

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $x, y-1, z$; (iii) $x, -y+2, z+1/2$.